

## Supplementary article data

# Cirrhosis is a risk factor for total hip arthroplasty for avascular necrosis

## A Danish nationwide cohort study

Thomas DELEURAN <sup>1,2</sup>, Søren OVERGAARD <sup>3</sup>, Hendrik VILSTRUP <sup>1</sup>, and Peter JEPSEN <sup>1,2</sup>

<sup>1</sup> Department of Hepatology and Gastroenterology and <sup>2</sup> Department of Clinical Epidemiology, Aarhus University Hospital, Aarhus; <sup>3</sup> Department of Orthopedic Surgery and Traumatology, Odense University Hospital, Institute of Clinical Research, University of Southern Denmark, Odense, Denmark.  
Correspondence: thomas.deleuran@clin.au.dk  
Submitted 2015-09-16. Accepted 2015-12-03.

Table 3. Alcohol-related disorders

	ICD-8	ICD-10
Alcohol-induced pseudo-Cushing's syndrome		D24.4
Alcoholic cardiomyopathy		I42.6
Alcoholic gastritis		K29.2
Alcoholic myopathy		G72.1
Alcoholic polyneuropathy		G62.1
Acute alcoholic pancreatitis		K85.2
Chronic alcoholic pancreatitis		K86.0
Degeneration of nervous system due to alcohol		G31.2
Ethanol poisoning		T51.0
Mental and behavioral disorders due to alcohol intake	291.xx, 303.xx	F10.0