

## Supplementary article data

# Muscular strength after total hip arthroplasty

## A prospective comparison of 3 surgical approaches

Siri B WINTHER<sup>1,2</sup>, Vigdis S HUSBY<sup>3</sup>, Olav A FOSS<sup>1</sup>, Tina S WIK<sup>1</sup>, Svein SVENNINGSEN<sup>4</sup>, Monika ENGDAL<sup>5</sup>, Kristin HAUGAN<sup>1</sup>, and Otto S HUSBY<sup>1</sup>

<sup>1</sup> Orthopaedic Research Centre, Orthopaedic Department, Trondheim University Hospital, Trondheim; <sup>2</sup> Department of Neuroscience, Norwegian University of Science and Technology, Trondheim; <sup>3</sup> Faculty of Nursing, Sør-Trøndelag University College, Trondheim; <sup>4</sup> Department of Orthopaedics, Sørlandet Hospital, Arendal; <sup>5</sup> Department of Physiotherapy, Trondheim University Hospital, Trondheim, Norway.  
Correspondence: siri.bjorgen@ntnu.no  
Submitted 2015-02-13. Accepted 2015-06-04.




Figure 1. Flow diagram showing the enrollment and inclusion process.




Figure 2. Set-up for the leg-press exercise.




Figure 3. Set-up for the abduction exercise.