

Supplementary article data

Increased migration of uncemented acetabular cups in female total hip arthroplasty patients with low systemic bone mineral density

A 2-year RSA and 8-year radiographic follow-up study of 34 patients

Sami FINNILÄ¹, Niko MORITZ¹, Erkki SVEDSTRÖM², Jessica J ALM¹, and Hannu T ARO¹

¹ Orthopaedic Research Unit, Turku University Hospital and University of Turku; ² Department of Diagnostic Radiology, Turku University Hospital, Turku, Finland.

Correspondence: hannu.aro@utu.fi

Submitted 2015-04-18. Accepted 2015-09-24.

Table 3. Evaluation of the stability of individual RSA markers and sufficiency of distribution at different time points of the postoperative follow-up. Values are median (range)

Segment	3 months	6 months	12 months	24 months
Mean error of rigid body fitting				
Bone markers	0.13 (0.04–0.33)	0.16 (0.05–0.31)	0.19 (0.04–0.36)	0.19 (0.04–0.33)
Implant markers	0.15 (0.04–0.30)	0.15 (0.01–0.29)	0.16 (0.04–0.37)	0.14 (0.04–0.36)
Condition number				
Bone markers	86 (33–180)	71 (33–179)	76 (33–182)	67 (33–183)
Implant markers	74 (34–117)	66 (31–94)	65 (34–91)	74 (36–113)